


Hellenic Sports Medicine
ΑΘΛΗΤΙΑΤΡΙΚΗ



www.sportsmedicinegreece.com

Τόμος 1, Τεύχος 1, Δεκέμβριος 2006
VOLUME 1 - ISSUE 1, DECEMBER 2006



HELLENIC SPORTS MEDICINE

Journal of the Sports Medicine
Association of Greece (S.M.A.G.)

Founder
K. Natsis

Publisher
K. Natsis
S.M.A.G. President

Property
Sports Medicine
Association of Greece
P.O. Box 1616, 54124, Thessaloniki

Printing House
PMP
(Paschalidis Medical Publications, Ltd)
14th, Tetrapoleos str.,
Athens, 115-27, Greece
Tel.: 0030-210-7789.125,
Fax: 0030-210-7759.421

Publishing Committee
P. Gigis
G. Godolias
N. Malliaropoulos
K. Natsis
Em. Papakostas
I. Terzidis

EDITORIAL BOARD

Director
K. Natsis

Members
N. Anastasopoulos
E. Asouhidou
I. Gigis
V. Karampatakis
K. Manavis
An. Beletsiotis
Em. Papakostas
N. Raikos
I. Terzidis

ΑΘΛΗΤΙΑΤΡΙΚΗ

Περιοδική Έκδοση της Αθλητιατρικής
Εταιρείας Ελλάδος (Α.Ε.Ε.)

Ιδρυτής
Κ. Νάτσης

Εκδότης
Κ. Νάτσης
Πρόεδρος Α.Ε.Ε.

Ιδιοκτησία
Αθλητιατρική
Εταιρεία Ελλάδος
Τ.Θ. 1616, 54124 Θεσσαλονίκη

Εκδοτικός Οίκος
Ιατρικές Εκδόσεις
Π.Χ Πασχαλίδης
Τετραπόλεως 14,
115 27 Αθήνα
Τηλ.: 0030-210.7789.125
Fax: 0030-210.7759.421

Εκδοτική Επιτροπή
Π. Γιγής
Γ. Γκοδόλιας
Ν. Μαλλιαρόπουλος
Κ. Νάτσης
Ε. Παπακώστας
Ι. Τερζίδης

ΣΥΝΤΑΚΤΙΚΗ ΕΠΙΤΡΟΠΗ

Διευθυντής
Κ. Νάτσης

Μέλη
Ν. Αναστασόπουλος
Ε. Ασουχίδου
Ι. Γιγής
Β. Καραμπατάκης
Κ. Μαναβής
Αν. Μπελετσιώτης
Εμ. Παπακώστας
Ν. Ράικος
Ι. Τερζίδης

Journal Secretariat

I. Terzidis
S. Levva

Γραμματεία Περιοδικού

I. Τερζίδης
Σ. Λέββα

Consulting Editors

K. Vasilikos
A. Geogoulis
I. Gigis
G. Gkioulekas
E. Tsoukali-Papadopoulou
D. Radopoulos
Th. Lialiaris
P. Nikolaou
An. Manthos
S. Papastergiou
J. Tsitouridis
An. Christodoulou
I. Christoforidis

Σύμβουλοι Έκδοσης

K. Βασιλικός
Α. Γεωργούλης
Π. Γιγής
Γ. Γιουλέκας
Ε. Τσούκαλη-Παπαδοπούλου
Δ. Ραδόπουλος
Θ. Λιαλιάρης
Π. Νικολάου
Α. Μάνθος
Σ. Παπαστεργίου
Ι. Τσιτουρίδης
Α. Χριστοδούλου
Ι. Χριστοφορίδης

International Editorial Board

H. J. Appell
N. Bachl
K. Christodoulakis
J. Duarte
G. Godolias
J. Koebke
U. Moebius
F. Pigozzi

Ετήσια Συνδρομή

Φυσικά Πρόσωπα: 30€

Φοιτητές: 10€

Ιδρύματα, Οργανισμοί κ.λ.π.: 40€

Διανέμεται Δωρεάν
στα Ταμειακώς Τακτοποιημένα Μέλη της Α.Ε.Ε.

Εγγραφές, Εμβάσματα (συνδρομών)-Διαφημίσεις

Αθλητιατρική Εταιρεία Ελλάδος


Τ.Θ. 1616, 54124 Θεσσαλονίκη

e-mail: natsis@med.auth.gr

Τραπεζικός Λογαριασμός:

Τράπεζα Eurobank,

Αριθμός: 0026.0139.32.0100519422



**SPORTS MEDICINE
ASSOCIATION OF GREECE**

Management Committee

President: K. Natsis

1st Vice President: P. Gigis

2nd Vice President: P. Nikolaou

Gen. Secretary: I. Terzidis

Ass. Secretary: N. Malliaropoulos

Treasurer: E. Papacostas

Library Supervisor: A. Porfiriadou

Members: N. Anastasopoulos

A. Beletsiotis

Ch. Bikos

S. Galitsanos

G. Godolias

K. Manavis

A. Papoutsidakis

N. Tzouroudis

**ΑΘΛΗΤΙΑΤΡΙΚΗ
ΕΤΑΙΡΕΙΑ ΕΛΛΑΔΟΣ**

Διοικητικό Συμβούλιο

Πρόεδρος: Κ. Νάτσης

Α' Αντιπρόεδρος: Π. Γιγής

Β' Αντιπρόεδρος: Π. Νικολάου

Γεν. Γραμματέας: Ι. Τερζίδης

Ειδ. Γραμματέας: Ν. Μαλλιαρόπουλος

Ταμίας: Ε. Παπακώστας

Εφ. Βιβλιοθήκης: Α. Πορφυριάδου

Μέλη: Ν. Αναστασόπουλος

Σ. Γαλιτσάνος

Γ. Γκοδόλιας

Κ. Μαναβής

Α. Μπελετσιώτης

Χ. Μπίκος

Α. Παπουτσιδάκης

Ν. Τζουρούδης



www.sportsmedicinegreece.com

Instructions for Authors

The "Hellenic Sports Medicine" is the official journal of the Sports Medicine Association of Greece. The journal publishes original papers, reviews, short communications and letters to the Editors from Greece or abroad. Manuscripts submitted to the journal must contain novel data on theoretical, clinical or experimental research or on practical applications in the field of sports medicine. No substantial part of the submission should have been published elsewhere. If a part of the submission has been published or presented on a congress, symposium, or national meeting proceeding, the reference for that publication and/or presentation should be given in the manuscript acknowledgement section. Submitted papers undergo peer reviewing by three independent reviewers.

CATEGORIES OF ARTICLES ACCEPTED FOR REVIEW

Review articles: Review articles on topics of broad interest are desirable. Authors who wish to submit an unsolicited review article should correspond with the Editors in Chief to determine the timeliness of the proposed review article. The correspondence should include an abstract and a complete outline of the proposed review article, including figures and tables (if possible). Review articles should not exceed 20 pages or 24,000 characters, including references and figures. Review articles are considered by the Editors and expert reviewers before a final decision regarding publication is made. Authors submitting review manuscripts should include a section describing the methods used for locating, selecting, extracting and synthesizing data. These methods should also be summarized in the abstract.

Original articles: Clinical, theoretical or experimental (basic or applied) research or practical applications in the field of Sports Medicine. Original articles should not exceed a total of 10 pages or 15,000 characters, including tables, figures and references. References should not exceed a total of 50. The form required include an abstract (no more than 250 words), a brief introduction where the purpose of the study should be referred, methods, statistical processing, results, discussion and references.

Rapid communications: Short manuscripts containing results of unusual scientific interest and importance, requiring rapid publication. Papers sub-

mitted for this accelerated process must require little or no revision. Accompanying the submission should be a letter from the authors stating why they feel the paper should be published as a rapid communication. The manuscripts must not exceed four journal pages (2,500 words including figures, tables and references). The format is the same as for full papers. References should not exceed 15.

Information articles: It is presented recent achievements in the field of Sports Medicine and they should not exceed six pages.

Proceedings, supplements and lectures of Sports Medicine congress.

Letters to the Editor: They are welcome and will be published if appropriate. Letters (maximum length 800 words) relating to material previously published in "Hellenic Sports Medicine" should be submitted within 6 months after publication of the material the letter is referring to.

STYLE OF THE MANUSCRIPTS

Manuscripts may be rejected without review on the basis of poor English or lack of conformity to stated standards of style. The text of observational and experimental articles is usually divided into sections with the headings: Introduction, Methods, Results and Discussion. Other types of articles, such as case reports, reviews and editorials, are likely to need other formats. Type or print on only one side of the paper. Use double-spacing throughout, including for the title page, abstract, text, acknowledgements, references, individual tables, and legends. There must be a space of 2.5 cm on each side of the paper.

First page: It should be referred the title. The title should be concise but informative. Names and addresses of the authors should not appear elsewhere in the main document, but in the bottom of the first page. The name of the clinic or the institution where the manuscript is coming from should also be referred in the first page. These data are entered separately.

Abstract: The abstract should be informative, self-explanatory without reference to the text of the manuscript. It should include essential significant results that support the conclusion of the work. It should also include the title and names of the authors.

Key-words should be 3-5 and representative of the subject studied or discussed.

Introduction: It should be comprehensible to the general reader, give a clear statement of the purpose of the paper and provide relevant context to support the basis for the paper and the significance of the work. Do not exhaustively review the literature or refer results from the work having done.

Materials & Methods: Provide sufficient information in the text or by reference to other work to permit the submitted work to be repeated without the need to communicate with the authors. Relevant validity and reliability data should be provided for critical methods. State the type of statistical tests used. Include the number of observations and the statistical findings when appropriate. Parametric and nonparametric statistics must be used as appropriate. When reporting experiments on human subjects, indicate whether the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation and with the Helsinki Declaration of 1975, as revised in 1983. Do not use patients' names, initials or hospital numbers, especially in illustrative material such as figures. When reporting experiments on animals, indicate whether the institution's or a national research council's guide for, or any national law on, the care and use of laboratory animals was followed.

Results: Should be presented precisely and should not contain material that is appropriate in the discussion. Units, quantities, and formulas should be expressed according to the Systeme Internationale (SI units). All measurements should be given in metric units.

Discussion: Emphasize the new and important aspects of the study and conclusions derived from the study.

Acknowledgements: These should be as brief as possible. Any grant that requires acknowledgement should be mentioned. The names of funding organizations should be written in full.

References: Titles of journals should be abbreviated according to the latest edition of Index Medicus. All authors should be named (do not use "et al."). Authors bear complete responsibility for the accuracy of the references. The references must be written according to Vancouver System. Number references (arabic numbers) consecutively in the order in which they are first mentioned in the text. Type references in the text by Arabic numbers as superscripts (above

the line of the text) and if at the end of a sentence after the period.

Only published or "in press" papers or books may be cited in the reference list. Information from manuscripts submitted but not yet accepted should be cited in the text as "unpublished observations" in parentheses. Personal communications should be listed in the text in parentheses. Published abstracts could be used as references only if the full text has not been published yet. Use of a large number of abstracts or non peer reviewed articles in the reference section will be grounds for rejection of the submission without review.

a) Journal article: Hubbard TJ, Hertel J. Mechanical contributions to chronic lateral ankle instability. *Sports Med.* 2006;36(3):26377.

b) Complete book: Hutson M.A. Sports Injuries. Recognitions and management. Oxford University Press, Oxford, 1990.

c) Chapter in a book: Phillips SJ, Whisnant JP. Hypertension and stroke. In: Leregh JH, Brenner BM, editors, Hypertension: pathophysiology, diagnosis, and management. 2nd ed. New York: Raven Press; 1995. p. 465-78.

d) Conference/Congress proceedings: Tocitu D, editor Enzymatic of the adaptative processes in High-Performance Sport. Proceedings of the 12th Balkan Sports Medicine Congress, 6th International Congress of the Sports Medicine Association of Greece, 3rd Hellenic-Cyprus Sports Medicine Congress 2002 March 21-24; Thessaloniki, Greece.

e) Dissertation: K.Natsis. Ultrastructural study of the skeletal muscle fibers after an experimental muscle atrophy [dissertation number 870]. Medical School, Aristotle University of Thessaloniki, 1993.

F) Journal article in electronic format: Christodoulou A, Terzidis i, Natsis k, Gigis i, Pournaras j. Soleus accessorius, an anomalous muscle in a young athlete: case report and analysis of the literature. *Br J Sports Med* 2004;38(6):e-38

Available from: URL:

<http://bjsm.bmj.com/cgi/content/full/38/6/e38>

Figures: Figures, illustrations, or halftones should be used when findings are best visually communicated. The use of photographs or equipment and experimental subjects should be avoided; good line drawings are more informative. Abbreviations used in the figure must be explained in the legend. Reference to the figure should be made in the text. Figures, illustrations or halftones must be sharp and high contrast. Uniform typographical setup (font style & size,

line thickness) of all figures in a paper is highly desirable. Images should be provided as .tif or .jpg files in a resolution of 300 dpi. Color figures, illustrations or halftones will not be published unless the author requires color in the publication. In this case the author will be charged with the additional cost of printing.

Tables: Tables should be used to communicate information that is hard to present visually. Results whose interpretation is more easily comprehended by knowing the means and SEM (or SD) may be presented in a table(s). Tables should be self-explanatory and bear a short title. Tables should be provided either as word or excel files.

Legends: Table and figure legends should be typed on separate sheet. A footnote to the table should explain all abbreviations used in the table.

SUBMITTING INFORMATION

- 1) Three full complimentary copies are supplied.
- 2) Every article should be followed by a cover letter citing the category of the article, that the article has not been published in the past in a Greek or international journal and that the authors demise the auctorial authorities to the Editorial Board. All the authors should sign the letter, which should not exceed the total of 800 words.
- 3) The article should be submitted necessarily in a "Word file" (CD or disk 3,5 A). Figures should submitted in tif or jpg files having a resolution at least 300 dpi.

GENERAL INFORMATION

After 6-8 weeks from receiving the article, the Editorial Board is going to inform the authors for accepting or not the manuscript.

Numbers one to nine should be referred fully written and up to ten arithmetically. Percentages should be referred always digitally, while a number in the beginning of a sentence should be fully written. Abbreviations should be written initially in parenthesis next to the full phrase.


Studies published in "Hellenic Sports Medicine" constitute literary property of the journal. Republishing partial or total is allowed only after written permission of the Editorial Board.

Papers must be sent to:

FOR THE JOURNAL HELLENIC SPORTS MEDICINE

Kostantinos Natsis, MD, PhD
Orthopaedic Surgeon
Pl. Ippodromiou . 17
P.C. 546 21 Thessaloniki
GREECE

email: natsis@med.auth.gr



Contents

- The arg577ter polymorphism in exon 16 of ACTN3 gene
in elite Greek field and track athletes 17**
Papadimitriou I.
- The effect of motor control physiotherapy after a hang glider's
surgically treated multiple trauma 20**
Tasheva R.
- Morphological and immunohistochemical study of the rat
myocardium after testosterone administration 24**
*Papamitou T., Dermentzopoulou-Theodoridou M., Barlagiannis D.,
Kotouza A., Kotanidou E., Manthos A.*
- Sport activity and development of osteoarthritis 29**
Iosifidis M.
- Injuries during Greek traditional dances 38**
Prantzidis I., Natsis K., Sofidis G., Kagioglou O., Prantzidis D.
- Rupture of anterior cruciate ligament and menstrual cycle
phase 43**
*Papastergiou P., Koukoulis N., Pappis G., Ziogas E.,
Dimitriadis Th., Parisis K.*
- Cranio-cerebral injuries during football games 51**
*Sirmos N., Kapoutzis N., Televantos A., Kapoutzis P., Chatzinasiou E.,
Sirmou E., Sirmos Ch.*

ABSTRACTS

The arg577ter polymorphism in exon 16 of ACTN3 gene in elite Greek field and track athletes

Papadimitriou I.

Abstract

The study of genetic influence in the making of an Olympic champion is still in its nascence, but this study provides some findings regarding the influence of genes on athletic performance. The objective of this study was to elucidate the genetic differences between elite Greek power and endurance track athletes by analyzing a mononucleotide polymorphism in exon 16 of the ACTN3 gene. The obtained results demonstrated that the genotype and allele frequencies of the top power athletes are statistically significant different from a representative random sample of the Greek population ($P=0.010$). It is worth mentioning that all the Olympic level sprinters participating in this study had at least one R allele in their genome. Thus, it is evident that the arg577ter polymorphism of ACTN3 gene is one of the various gene polymorphisms determining the levels of power-sprint ability of the athletes.

Key words: Athletes, arg577ter polymorphism, ACTN3 gene

The effect of motor control physiotherapy after a hang glider's surgically treated multiple trauma

Tasheva R.

Abstract

Hang gliding is one of the most dangerous sport activities because of the risk of highly-energetic traumas including acetabular fractures. Modern trends in treating complex acetabular fractures are directed at applying a surgical method securing anatomical reduction, stable internal fixation and a possibility of early physiotherapy. The precise implementation of motor tasks in physiotherapy after the adequate choice and applied of surgical approaches leads to a faster overcoming of the negative phenomena from movement disorders and to return to sport activity.

The purpose of the study is to present physical therapy program after surgically treated complex trauma caused by hang gliding.

Key Words: hang gliding, motor control, physiotherapy.

Morphological and immunohistochemical study of the rat myocardium after testosterone administration

Papamitsou T., Dermentzopoulou-Theodoridou M., Barlagiannis D., Kotouza A., Kotanidou E., Manthos A.

Abstract

The principal androgen in male and female is testosterone, which along with its metabolite dihydrotestosterone, exerts anabolic and androgenic effects through the androgen receptor. Non genomic effects are also well documented. Exogenous anabolic androgenic steroids and their influence on the performance and function of the heart, muscle, liver and other tissues have gained great interest. However, the usefulness of exogenous steroids in the myocardium has not been documented. The aim of this study is to demonstrate the ultrastructural alterations caused in the rat heart when testosterone was administered in rats, as well as to explore immunohistochemically any changes in the expression of the androgen receptor. Two groups of adult male Wistar rats were used. The first group received an everyday intramuscular dose of 12mg testosterone enanthate for 20 days, while the second group received placebo via the same route and for the same period of time. All rats were anaesthetized and sacrificed after the 20th day and tissue samples from the myocardium were received and processed for

transmission electron microscopy and immunohistochemistry using specific monoclonal antibodies against the androgen receptor. Significant ultrastructural alterations were observed in the myocytes. We report pathological appearance of the sarcomeres' structure including disorganization of the Z line. Mitochondrial oedema and enlargement of the mitochondria were abundantly present. Mitochondrial fusion was significant as was the existence of glycogen particles in between the sarcomeres. Immunohistochemistry showed the expression of the androgen receptor on the control group. When testosterone was administered, the staining for androgen receptor in the myocytes was more dense which was considered as an increase in the expression of the androgen receptor. Exogenous testosterone produces pathological alterations in the structure of myocytes in the rat heart. The androgen receptor increases its expression and may play a role in the observed ultrastructural changes. These findings are in agreement with studies that report the androgen receptor mediated cardiac hypertrophy caused by steroid abuse.

Key words: testosterone, morphological study, immunohistochemical study, myocardium, rat

Sport activity and development of osteoarthritis

Iosifidis M.

A b s t r a c t

Elite athletes' sports activities cause increased loads in their joints. Injuries of lower extremities increase the possibility of osteoarthritis. However, if the high impact and torsional loading and also the repeated low magnitude loading generated during intensive sport activity, cause cartilage degeneration is not clear yet. In our study we investigated the possible development of lower extremities osteoarthritis in former Greek elite athletes.

We studied 218 former elite athletes (soccer players, skiers, volleyball players, martial arts athletes, track and fields athletes, basketball players/ range of ages 40-84 years, mean 50.1 years SD=8.5). The control group was 181 males (range of ages 40-77 years, mean 50.7 years, SD=10) who didn't have systematic sport activity. The participants in the study had not diagnosed or/and operated for significant lower limb injury. Osteoarthritis was recorded through questionnaire, clinical examination and radiographic evaluation.

After adjusting the age, height, weight and body mass index (BMI), the statistical analysis (SPSS, independent samples t-test, z-test, CI=95%) showed no difference in the osteoarthritis incidence in former elite Greek athletes compared with the general population ($p>0,05$). However, the intensive sport activity was the major risk factor for the athletes, when the control group's body weight, occupation, and age were the main risk factors in the development of OA. Radiographic examination showed that the images of the former athletes presented increased osteoarthritic signs (osteophytes, joint line sclerosis, joint narrowing) in comparison with the control group, without, however, clinical signs of OA (pain, impaired ROM) ($p<0,01$).

Key words: Sport Activity - Articular Cartilage Degeneration – Osteoarthritis

Injuries during Greek traditional dances

Prantzidis I., Natsis K., Sofidis G., Kagioglou O., Prantzidis D.

A b s t r a c t

With the urbanization and the industrialization of Greek community, the Greek Traditional Dance (G.T.D.) is breaking away from its natural environment and is being transferred to the city as a spectacle. The consequence of this was a gradual extension in the teaching of traditional dance in schools and dancing societies. As a result there is a gradual increase of people who are involved with traditional dances, the establishment of dancing societies and the creation of a new occupation (a teacher of G.T.D.). The natural aftermath of this organized and methodical occupation with G.T.D. was the increase of the percentage of injuries in the people who are involved with it.

The purpose of this study was to register the kind and the frequency of dancer's lesions. Many confidential questionnaires were distributed and were filled namelessly by dancers from different regions of Greece. The specimen of the study was 485 individuals (202 men and 283 women), all members of a dancing society, with an average of age 27.68 years. The average of the occupation with dance was 9,01 years. From these people 77 persons were injured (15,87%), 43 were men (55,84%) and 34 women (44,15%). From the 77 individuals, 74 reported the part of the body which was injured, while 3 persons didn't. 71 dancers had an injury in one part of the body and 4 in two parts. From 78 injuries, 31 related to the ankle joint (39,74%), 23 to the knee joint (29,49%), 6 to the quadriceps muscle (7,69%), 4 to Achille tendon (5,13%), 3 to lumbar spine region (3,85%), 5 to foot (6,41%), 2 to tibia (2,56%), 2 to gastrocnemius muscle (2,56%), 1 to biceps femoris muscle (1,28%) and 1 to thumb. From 77 persons, 73 (94,8) visited a doctor for the diagnosis of the injury while 4 (5,20%) did not. From the lesions which were diagnosed 56 (71,79%) were characterized as acute injuries and 22 (28,21%) as chronic.

To 51 individuals (65,38%) a conservative treatment without medicine was recommended. A conservative treatment with medicine in 18 persons (2,08%) and in 9 dancers (11,54%) a surgery.

Key words: Greek traditional dances, injuries

Rupture of anterior cruciate ligament and menstrual cycle phase

Papastergiou P., Koukoulis N., Pappis G., Ziogas E., Dimitriadis Th., Parisis K.

Abstract

Purpose: Evaluation of a probable correlation between the anterior cruciate ligament rupture and a specific menstrual cycle phase in athletes.

Material and Method: Study of 42 female athletes of different sports who underwent an anterior cruciate ligament reconstruction, could define the day of menstrual cycle accurately when the knee injury sustained, weren't using oral contraceptives and had regular menstrual cycle of 28 days.

Results: From our study of 42 cases we found that 28 of women were injured during the follicular phase and 14 of women were injured during the luteal phase of their menstrual cycle.

Conclusion: There is correlation between the anterior cruciate ligament rupture and the menstrual cycle phase. Female athletes demonstrate higher than expected anterior cruciate ligament rupture rate during the follicular phase of menstrual cycle, and lower during the luteal phase of menstrual cycle (p=0.05).

Key words: Anterior Cruciate Ligament (ACL), Menstrual Cycle Phase.

Craniocerebral injuries during football games

Sirmos N., Kapoutzis N., Televantos A., Kapoutzis P., Chatzinasiou E., Sirmou E., Sirmos Ch.

Abstract

During Football games many accidents between the players may happen.

In the world of sports, soccer is unique becomes of the purpose for use of the unprotected head for controlling and advancing the ball.

Head injuries (craniocerebral, craniofacial) can be a result of contact of the head with another head (or other body parts), ground, goal post, other unknown objects or even the ball, almost never the less is a safe sport. We present our experience for emergency treatment in our surgical department in a peripheral hospital and Health Center during the period 2002 - 2005.

Key words: football, craniocerebral - craniofacial injuries - accident in football players.