

Hellenic Sports Medicine
ΑΘΛΗΤΙΑΤΡΙΚΗ



www.sportsmedicinegreece.com

Τόμος 2, Τεύχος 1, Μάιος 2007
VOLUME 2 - ISSUE 1, May 2007



HELLENIC SPORTS MEDICINE

Journal of the Sports Medicine
Association of Greece (S.M.A.G.)

Founder

K. Natsis

Publisher

K. Natsis

S.M.A.G. President

Property

Sports Medicine

Association of Greece

P.O. Box 1616, 54124, Thessaloniki

Printing House

PMP

(Paschalidis Medical Publications, Ltd)

14th, Tetrapoleos str.,

Athens, 115-27, Greece

Tel.: 0030-210-7789.125,

Fax: 0030-210-7759.421

Publishing Committee

P. Gigis

G. Godolias

N. Malliaropoulos

K. Natsis

Em. Papakostas

I. Terzidis

EDITORIAL BOARD

Director

K. Natsis

Members

N. Anastasopoulos

E. Asouhidou

I. Gigis

V. Karampatakis

K. Manavis

An. Beletsiotis

Em. Papakostas

N. Raikos

I. Terzidis

ΑΘΛΗΤΙΑΤΡΙΚΗ

Περιοδική Έκδοση της Αθλητιατρικής
Εταιρείας Ελλάδος (Α.Ε.Ε.)

Ιδρυτής

Κ. Νάτσης

Εκδότης

Κ. Νάτσης

Πρόεδρος Α.Ε.Ε.

Ιδιοκτησία

Αθλητιατρική

Εταιρεία Ελλάδος

Τ.Θ. 1616, 54124 Θεσσαλονίκη

Εκδοτικός Οίκος

Ιατρικές Εκδόσεις

Π.Χ Πασχαλίδης

Τετραπόλεως 14,

115 27 Αθήνα

Τηλ.: 0030-210.7789.125

Fax: 0030-210.7759.421

Εκδοτική Επιτροπή

Π. Γιγής

Γ. Γκοδόλιας

Ν. Μαλλιαρόπουλος

Κ. Νάτσης

Ε. Παπακώστας

Ι. Τερζίδης

ΣΥΝΤΑΚΤΙΚΗ ΕΠΙΤΡΟΠΗ

Διευθυντής

Κ. Νάτσης

Μέλη

Ν. Αναστασόπουλος

Ε. Ασουχίδου

Ι. Γιγής

Β. Καραμπατάκης

Κ. Μαναβής

Αν. Μπελετσιώτης

Εμ. Παπακώστας

Ν. Ράικος

Ι. Τερζίδης

Journal Secretariat

I. Terzidis
A. Adamopoulou

Γραμματεία Περιοδικού

I. Τερζίδης
Α. Αδαμοπούλου

Consulting Editors - Reviewers

M. Albani
K. Vasilikos
A. Georgoulis
P. Gigis
G. Gioulekas
G. Kapetanos
T. Lialiaris
A. Manthos
P. Baltopoulos
P. Nikolaou
N. Dombros
S. Papastergiou
D. Radopoulos
P. Skandalakis
P. Tsikaras
I. Tsitouridis
M. Tsolaki
E. Tsoukali-Papadopoulou
A. Christodoulou
I. Christoforidis

Σύμβουλοι Έκδοσης - Κριτές

M. Αλμπάνη
Κ. Βασιλικός
Α. Γεωργούλης
Π. Γιγής
Γ. Γιουλέκας
Γ. Καπετάνος
Θ. Λιαλιάρης
Α. Μάνθος
Π. Μπαλτόπουλος
Π. Νικολάου
Ν. Ντόμπρος
Σ. Παπαστεργίου
Δ. Ραδόπουλος
Π. Σκανδαλάκης
Π. Τσικάρας
Ι. Τσιτουρίδης
Μ. Τσολάκη
Ε. Τσούκαλη-Παπαδοπούλου
Α. Χριστοδούλου
Ι. Χριστοφορίδης

International Editorial Board

H. J. Appell
N. Bachl
K. Christodoulakis
J. Duarte
G. Godolias
A. Ionescu
J. Koebke
U. Moebius
F. Pigozzi

Ετήσια Συνδρομή

Φυσικά Πρόσωπα: 30€
Φοιτητές: 10€
Ιδρύματα, Οργανισμοί κ.λ.π.: 40€

Διανέμεται Δωρεάν στα Ταμειακώς
Τακτοποιημένα Μέλη της Α.Ε.Ε.

Εγγραφές, Εμβάσματα (συνδρομών)-Διαφημίσεις

Αθλητιατρική Εταιρεία Ελλάδος
Τ.Θ. 1616, 54124 Θεσσαλονίκη
e-mail: natsis@med.auth.gr
Τραπεζικός Λογαριασμός:
Τράπεζα Eurobank,
Αριθμός: 0026.0139.32.0100519422



**SPORTS MEDICINE
ASSOCIATION OF GREECE**

Management Committee

President: K. Natsis

1st Vice President: P. Gigis

2nd Vice President: P. Nikolaou

Gen. Secretary: I. Terzidis

Ass. Secretary: N. Malliaropoulos

Treasurer: E. Papacostas

Library Supervisor: A. Porfiriadou

Members: N. Anastasopoulos

A. Beletsiotis

Ch. Bikos

S. Galitsanos

G. Godolias

K. Manavis

A. Papoutsidakis

N. Tzouroudis

**ΑΘΛΗΤΙΑΤΡΙΚΗ
ΕΤΑΙΡΕΙΑ ΕΛΛΑΔΟΣ**

Διοικητικό Συμβούλιο

Πρόεδρος: Κ. Νάτσης

Α' Αντιπρόεδρος: Π. Γιγής

Β' Αντιπρόεδρος: Π. Νικολάου

Γεν. Γραμματέας: Ι. Τερζίδης

Ειδ. Γραμματέας: Ν. Μαλλιαρόπουλος

Ταμίας: Ε. Παπακώστας

Εφ. Βιβλιοθήκης: Α. Πορφυριάδου

Μέλη: Ν. Αναστασόπουλος

Σ. Γαλιτσάνος

Γ. Γκοδόλιας

Κ. Μαναβής

Α. Μπελετσιώτης

Χ. Μπίκος

Α. Παπουτσιδάκης

Ν. Τζουρούδης



www.sportsmedicinegreece.com

Instructions for Authors

The "Hellenic Sports Medicine" is the official journal of the Sports Medicine Association of Greece. The journal publishes original papers, reviews, short communications and letters to the Editors from Greece or abroad. Manuscripts submitted to the journal must contain novel data on theoretical, clinical or experimental research or on practical applications in the field of sports medicine. No substantial part of the submission should have been published elsewhere. If a part of the submission has been published or presented on a congress, symposium, or national meeting proceeding, the reference for that publication and/or presentation should be given in the manuscript acknowledgement section. Submitted papers undergo peer reviewing by three independent reviewers.

CATEGORIES OF ARTICLES ACCEPTED FOR REVIEW

Review articles: Review articles on topics of broad interest are desirable. Authors who wish to submit an unsolicited review article should correspond with the Editors in Chief to determine the timeliness of the proposed review article. The correspondence should include an abstract and a complete outline of the proposed review article, including figures and tables (if possible). Review articles should not exceed 20 pages or 24,000 characters, including references and figures. Review articles are considered by the Editors and expert reviewers before a final decision regarding publication is made. Authors submitting review manuscripts should include a section describing the methods used for locating, selecting, extracting and synthesizing data. These methods should also be summarized in the abstract.

Original articles: Clinical, theoretical or experimental (basic or applied) research or practical applications in the field of Sports Medicine. Original articles should not exceed a total of 10 pages or 15,000 characters, including tables, figures and references. References should not exceed a total of 50. The form required include an abstract (no more than 250 words), a brief introduction where the purpose of the study should be referred, methods, statistical processing, results, discussion and references.

Rapid communications: Short manuscripts containing results of unusual scientific interest and importance, requiring rapid publication. Papers sub-

mitted for this accelerated process must require little or no revision. Accompanying the submission should be a letter from the authors stating why they feel the paper should be published as a rapid communication. The manuscripts must not exceed four journal pages (2,500 words including figures, tables and references). The format is the same as for full papers. References should not exceed 15.

Information articles: It is presented recent achievements in the field of Sports Medicine and they should not exceed six pages.

Proceedings, supplements and lectures of Sports Medicine congress.

Letters to the Editor: They are welcome and will be published if appropriate. Letters (maximum length 800 words) relating to material previously published in "Hellenic Sports Medicine" should be submitted within 6 months after publication of the material the letter is referring to.

STYLE OF THE MANUSCRIPTS

Manuscripts may be rejected without review on the basis of poor English or lack of conformity to stated standards of style. The text of observational and experimental articles is usually divided into sections with the headings: Introduction, Methods, Results and Discussion. Other types of articles, such as case reports, reviews and editorials, are likely to need other formats. Type or print on only one side of the paper. Use double-spacing throughout, including for the title page, abstract, text, acknowledgements, references, individual tables, and legends. There must be a space of 2.5 cm on each side of the paper.

First page: It should be referred the title. The title should be concise but informative. Names and addresses of the authors should not appear elsewhere in the main document, but in the bottom of the first page. The name of the clinic or the institution where the manuscript is coming from should also be referred in the first page. These data are entered separately.

Abstract: The abstract should be informative, self-explanatory without reference to the text of the manuscript. It should include essential significant results that support the conclusion of the work. It should also include the title and names of the authors.

Key-words should be 3-5 and representative of the subject studied or discussed.

Introduction: It should be comprehensible to the general reader, give a clear statement of the purpose of the paper and provide relevant context to support the basis for the paper and the significance of the work. Do not exhaustively review the literature or refer results from the work having done.

Materials & Methods: Provide sufficient information in the text or by reference to other work to permit the submitted work to be repeated without the need to communicate with the authors. Relevant validity and reliability data should be provided for critical methods. State the type of statistical tests used. Include the number of observations and the statistical findings when appropriate. Parametric and nonparametric statistics must be used as appropriate. When reporting experiments on human subjects, indicate whether the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation and with the Helsinki Declaration of 1975, as revised in 1983. Do not use patients' names, initials or hospital numbers, especially in illustrative material such as figures. When reporting experiments on animals, indicate whether the institution's or a national research council's guide for, or any national law on, the care and use of laboratory animals was followed.

Results: Should be presented precisely and should not contain material that is appropriate in the discussion. Units, quantities, and formulas should be expressed according to the Systeme Internationale (SI units). All measurements should be given in metric units.

Discussion: Emphasize the new and important aspects of the study and conclusions derived from the study.

Acknowledgements: These should be as brief as possible. Any grant that requires acknowledgement should be mentioned. The names of funding organizations should be written in full.

References: Titles of journals should be abbreviated according to the latest edition of Index Medicus. All authors should be named (do not use "et al."). Authors bear complete responsibility for the accuracy of the references. The references must be written according to Vancouver System. Number references (arabic numbers) consecutively in the order in which they are first mentioned in the text. Type references in the text by Arabic numbers as superscripts (above

the line of the text) and if at the end of a sentence after the period.

Only published or "in press" papers or books may be cited in the reference list. Information from manuscripts submitted but not yet accepted should be cited in the text as "unpublished observations" in parentheses. Personal communications should be listed in the text in parentheses. Published abstracts could be used as references only if the full text has not been published yet. Use of a large number of abstracts or non peer reviewed articles in the reference section will be grounds for rejection of the submission without review.

a) Journal article: Hubbard TJ, Hertel J. Mechanical contributions to chronic lateral ankle instability. *Sports Med.* 2006;36(3):26377.

b) Complete book: Hutson M.A. Sports Injuries. Recognitions and management. Oxford University Press, Oxford, 1990.

c) Chapter in a book: Phillips SJ, Whisnant JP. Hypertension and stroke. In: Leregh JH, Brenner BM, editors, Hypertension: pathophysiology, diagnosis, and management. 2nd ed. New York: Raven Press; 1995. p. 465-78.

d) Conference/Congress proceedings: Tocitu D, editor Enzymatic of the adaptative processes in High-Performance Sport. Proceedings of the 12th Balkan Sports Medicine Congress, 6th International Congress of the Sports Medicine Association of Greece, 3rd Hellenic-Cyprus Sports Medicine Congress 2002 March 21-24; Thessaloniki, Greece.

e) Dissertation: K.Natsis. Ultrastructural study of the skeletal muscle fibers after an experimental muscle atrophy [dissertation number 870]. Medical School, Aristotle University of Thessaloniki, 1993.

F) Journal article in electronic format: Christodoulou A, Terzidis i, Natsis k, Gigis i, Pournaras j. Soleus accessorius, an anomalous muscle in a young athlete: case report and analysis of the literature. *Br J Sports Med* 2004;38(6):e-38

Available from: URL:

<http://bjsm.bmj.com/cgi/content/full/38/6/e38>

Figures: Figures, illustrations, or halftones should be used when findings are best visually communicated. The use of photographs or equipment and experimental subjects should be avoided; good line drawings are more informative. Abbreviations used in the figure must be explained in the legend. Reference to the figure should be made in the text. Figures, illustrations or halftones must be sharp and high contrast. Uniform typographical setup (font style & size,

line thickness) of all figures in a paper is highly desirable. Images should be provided as .tif or .jpg files in a resolution of 300 dpi. Color figures, illustrations or halftones will not be published unless the author requires color in the publication. In this case the author will be charged with the additional cost of printing.

Tables: Tables should be used to communicate information that is hard to present visually. Results whose interpretation is more easily comprehended by knowing the means and SEM (or SD) may be presented in a table(s). Tables should be self-explanatory and bear a short title. Tables should be provided either as word or excel files.

Legends: Table and figure legends should be typed on separate sheet. A footnote to the table should explain all abbreviations used in the table.

SUBMITTING INFORMATION

- 1) Three full complimentary copies are supplied.
- 2) Every article should be followed by a cover letter citing the category of the article, that the article has not been published in the past in a Greek or international journal and that the authors demise the auctorial authorities to the Editorial Board. All the authors should sign the letter, which should not exceed the total of 800 words.
- 3) The article should be submitted necessarily in a "Word file" (CD or disk 3,5 A). Figures should submitted in tif or jpg files having a resolution at least 300 dpi.

GENERAL INFORMATION

After 6-8 weeks from receiving the article, the Editorial Board is going to inform the authors for accepting or not the manuscript.

Numbers one to nine should be referred fully written and up to ten arithmetically. Percentages should be referred always digitally, while a number in the beginning of a sentence should be fully written. Abbreviations should be written initially in parenthesis next to the full phrase.

Studies published in "Hellenic Sports Medicine" constitute literary property of the journal. Republishing partial or total is allowed only after written permission of the Editorial Board.

Papers must be sent to:

FOR THE JOURNAL HELLENIC SPORTS MEDICINE

Kostantinos Natsis, MD, PhD
Orthopaedic Surgeon
Pl. Ippodromiou . 17
P.C. 546 21 Thessaloniki
GREECE

email: natsis@med.auth.gr



Contents

- Examination of obesity degree in relation to physical skills on Elementary School pupils 17**
Mavrovouniotis F., Arsenopoulou M., Argiriadou Eir.
- Are there any distinct blood flow patterns in the thigh and the calf of athletes, healthy volunteers and patients with claudication at rest, during, and after exercise? 26**
Angelides N., Konstandinou Ch., Costeas A. , Christofides N.
- Body composition comparison between athletes, anorexia nervosa patients and recovered former anorexia nervosa patients, using D.E.X.A. 32**
Grammatikopoulou M.G., Stewart A.D., Zakas A., Tzetzis G., Grammatikopoulos G.
- Physiotherapy in conservative treatment of spondylolysis in athletes 39**
Tasheva R., Ganchev D.
- Liquid-liquid extraction of amphetamine from biological samples of blood and urine 43**
Raikos N., Pouliopoulos A., Spagou K., Natsis K., Psaroulis D., Tsoukali E.
- The frequency of injuries in kickboxing 49**
Kotsikas G., Kitsios A., Sykaras E., Teflioudis A.

Examination of obesity degree in relation to physical skills on Elementary School pupils

Mavrovouniotis F., Arsenopoulou M., Argiriadou Eir.

Abstract

The purpose of the present study was the investigation of obesity degree in relation to sport tests on Elementary School pupils. For this aim 271 pupils of the six grades of the primary school were examined (148 boys and 123 girls). Measurements of height, body mass, waist and hip perimeter were performed. Moreover, tests in long jump, vertical jump and trunk flexion were carried out. Body Mass Index was used for the evaluation of the degree of overweight and obesity, while waist perimeter was used for the evaluation of the degree of central obesity. From data statistical analysis it was found out high percentages of central obesity in all ages that fluctuated from 6.9% up to 38.9%. In addition, high percentages of overweight and obese children were observed, both boys (38.66% and 13.31%, respectively) and girls (39.96% and 17.60%, respectively). Important is the fact that overweight and obese children come significantly behind in the long jump ($p < 0.001$) and vertical jump ($p < 0.01$ and $p < 0.001$) performance, against children of normal weight. This decreased physical robustness and muscle strength of overweight and obese children distinguishes inactivity and decreased participation in physical activities, as well as tendency for lower physical fitness. Consequently, obesity prevention in childhood is imposed, with ways mainly the increase of physical activity that involves inactivity reduction and television and videogames employment, but also the controlled diet, as the most effective strategy for obesity control.

Key words: central obesity, physical activity, diet.

Are there any distinct blood flow patterns in the thigh and the calf of athletes, healthy volunteers and patients with claudication at rest, during, and after exercise?

Angelides N., Konstandinou Ch., Costeas A., Christofides N.

Abstract

The purpose of this study was to investigate whether there are distinct patterns of flow in the thigh and calf of athletes, normal individuals and patients with peripheral arterial disease, under resting conditions, during walking and running on the treadmill, as well as during the post-exercise period. For this purpose simultaneous ^{99m}Tc clearance in the thigh and in the calf was used at rest, during and after exercise. The study was based on the principle that the clearance of the radioactive isotope was proportional to blood flow in the thigh and in the calf at any given time. All individuals rested before starting the test. Then, forty μCi of ^{99m}Tc diluted in 0.2 ml of saline were injected into the thickest part of the quadriceps and gastrocnemius muscles. Two light probes were strapped on the skin over the points of injections. These probes were connected to two isotope localization monitors, and the output from the detectors was recorded continuously on a pen recorder. Clearance curves were recorded for three minutes at rest. Then, all individuals walked on a horizontal treadmill at 4.5 Km per minute for three minutes or until claudication occurred. All normal volunteers, athletes and non-athletes, were also asked to run on the treadmill at 9 Km per minute for another three minutes. Clearance curves were obtained at rest and during exercise. Finally clearance curves

were recorded during the post-exercise resting period for ten minutes. The percentage clearance per minute (T) at any minute, was obtained from the following equation (1):

$$T = \frac{f(t) - f(t+1)}{f(t) + f(t+1)} \times 200 \quad (\text{equation 1})$$

where $f(t)$ was the radioactivity measured at time t and $f(t+1)$ was the radioactivity measured one minute latter. The radioactivity cleared during one minute was $f(t) - f(t+1)$ while the mean radioactivity during the same minute was $f(t) + f(t+1) / 2$ and therefore the percentage radioactivity cleared per minute was given by equation 1. Athletes (sprinters), as well as non-athletes individuals with no sign of peripheral arterial disease and patients with claudication were included in the study. The results demonstrated that although muscle blood flow in the thigh and calf in normal individuals were comparable in athletes and non-athletes, the increase of flow was greater both in the thigh and in the calf of athletes during and after exercise, than in non-athletes individuals. The faster clearance of the radioactive isotope in the thigh and calf of athletes can only be explained if we take into consideration the well developed quadriceps and gastrocnemius muscles of these athletes. In patients with peripheral arterial occlusive disease the pattern of clearance during and after exercise is not the same. In limbs with proximal occlusion the result suggested that the blood flow pattern in the thigh and in the calf is the same as in normal limbs, but was "throttled" by the proximal occlusion so that blood flow was insufficient to permit normal hyperaemia. In limbs with distal occlusion, the hyperaemia in the thigh was normal, but flow in the calf was diminished immediately after exercise when there was maximum vasodilatation in the muscle bed of this area, but later showed a delayed hyperaemia when the hyperaemia in the thigh subsided. The only conceivable explanation of the diminished flow in the calf immediately after exercise is a fall in pressure in the axial vessels distal to the occlusion. This in turn is the consequence of a similar fall in the pressure gradient across the collateral circulation. The calf is deprived and can only begin to receive blood when the hyperaemia in the thigh subsides. This then permits hyperaemia in the calf where maximum dilatation is still present. In conclusion, there are distinct patterns of changes in flow during and after exercise, in athletes, in non-athletes normal individuals and in patients with peripheral arterial disease, which can be determined by ^{99m}Tc muscle clearance. These changes are so characteristic that they can be used for research and for diagnostic purposes.

Key words: blood flow quadriceps gastrocnemius isotope clearance during exercise.

Body composition comparison between athletes, anorexia nervosa patients and recovered former anorexia nervosa patients, using D.E.X.A..

Grammatikopoulou M.G., Stewart A.D., Zakas A., Tzetzis G., Grammatikopoulos G.

Abstract

The aim of this study was to assess total and regional body composition differences of female eating disorders patients and recovered eating disorders patients compared to athletes and a controls group.

Subjects formed 4 groups, 28 current anorexics, 32 recovered anorexics, 30 athletes and 30 controls, all Caucasian females. Dual-energy x-ray absorptiometry was used to examine total and regional body composition.

Results showed a similar percentage body composition for athletes and anorexics ($p=1$ for %lean, %fat, and %Bone Mineral Content). Athletes and anorexics also presented similar ($p=1$) % leg and torso fat whereas the RA group differed significantly for the same body regions, compared to the athletes ($p<0.05$). Controls had significantly different %torso fat compared to the patients and the athletes and different %leg fat compared to the athletes only.

Anorexics appeared to have similar body components of lean, fat and BMC to those of the athlete population. These similarities suggest that an unhealthy pattern for body image and appearance exists in the female athlete population.

Key words: eating disorders, body fat, anthropometry, bone mineral content.

Physiotherapy in conservative treatment of spondylolysis in athletes

Tasheva R., Ganchev D.

Abstract

The aim of this study is to develop the physiotherapy program in conservative treatment of spondylolysis in athletes.

Between December 2000 and December 2005 we observed 12 athletes with spondylolysis with mean age 17,6 (15-22) years. There were 8 female and 4 males. Average follow-up term is 11 (5-36) months. The physiotherapy program taught the patients to maintain the pelvis tilt and to restore the muscle balance.

The results were assayed on the ground of subjective (pain, discomfort, and neurological symptoms) and functional objective (low back and hip range of motion, extending the big toe, and athletic activity) criteria. Regarding the subjective criteria there are 2 (16,67%) patients with mild pain, discomfort and neurological symptoms. Two patients have some weakness in extending their big toe. All patients are with restore of low back and hip range of motion and only 1 (8, 3%) of them didn't feel comfortable during sports activity.

The combination of conservative treatments such as bracing, static and dynamic physiotherapeutic program, and lifestyle changes can control the condition of the athletes with spondylolysis. It is important to know that any therapeutic approach must take into account that spondylolysis means that there is a potentially unstable area of the spine, so the competency of the treating spine specialist is very important consideration.

Key words: athletes, conservative treatment, physiotherapy program, spondylolysis.

Liquid-Liquid extraction of amphetamine from biological samples of blood and urine

Raikos N., Pouliopoulos A., Spagou K., Natsis K., Psaroulis D., Tsoukali E.

Abstract

Amphetamine has been extensively used therapeutically in the past. However, its central stimulating actions have led to extensive abuse with attendant social and psychiatric problems. The objective of this study was the optimization of amphetamine extraction from biological samples by liquid-liquid extraction. The recovery of amphetamines was studied using a variety of 5 solvents (dichloromethane, 1-chlorobutan, ethyl acetate, di-ethylether, chloroform) as well as a system of solvents (chloroform: ethyl acetate: ethanol, 3:1:1 v/v) named as alkaline B. The extracting solvent was further examined for the appropriate pH value and quantity with the maximum recovery. After the optimization of the method, the recovery of amphetamines using

biological samples was studied. The determination of amphetamine was accomplished by gas chromatography equipped with a flame ionization detector (FID), after derivatization with heptafluorobutyric anhydride (HFBA).

Key words: amphetamine, liquid-liquid extraction, gas chromatography.

The frequency of injuries in Kickboxing

Kotsikas G., Kitsios A., Sykaras E., Teflioudis A.

Abstract

OBJECTIVES: This study was prepared with the cooperation of coaches and athletes and its purpose is to show the frequency of injuries in kickboxing. A valid questionnaire was used and the interviews were taken at the place where the athletes work - out.

METHODS: A total of 121 athletes participated in the study, 99 of them were males (81,8%) and 22 females (18,2%) (Graph 1). The mean age for males was the 22,90 years and the mean age for the females was the 20,27years (Chart 1). Out of the 121 athletes, 112 (93%) of them were amateurs and 9 (7%) were professionals (Graph 2).

RESULTS: The most common body region injured were the lower extremities 14%, followed by injuries to the wrist 9,9%. Injuries to the foot and thigh were 8,3%. Injuries to the neck region 5,8%, to the elbow region 3,3%. Injuries to the knee and hand were 3,3%, followed by injuries to the spine 1,7%, to the pelvic region and

the head 0,8% (Chart 2). The type of injury in most of the cases was identified as strain 55%, followed by

sprains 20% and fractures 5%, an 8% of them was a combination of different injuries (Graph 5).

Conclusion: Kickboxing is a safe sport, but further research into injury patterns in different styles of kickboxing and the mechanism of injury occurrence and also in relation with similar contact sports is required.

Key words: frequency of injuries, type of injuries, different age groups, kickboxing.